

JOURNAL OF DIFFERENTIAL EQUATIONS 41, 369-374 (1981)

An Anti-maximum Principle for Linear Elliptic Equations with an Indefinite Weight Function

PETER HESS

Mathematisches Institut, Universität Zurich, Zurich, Switzerland

Received July 8, 1980

1. INTRODUCTION AND STATEMENT OF THE RESULT

In this note we prove an anti-maximum principle for the Dirichlet problem

$$\begin{aligned}\mathcal{L}u - \lambda mu &= h && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega,\end{aligned}\tag{1}$$

in the bounded domain $\Omega \subset \mathbb{R}^N$ ($N \geq 1$) with smooth boundary $\partial\Omega$. By \mathcal{L} :

$$\mathcal{L}u = - \sum_{j,k=1}^N a_{jk} \frac{\partial^2 u}{\partial x_j \partial x_k} + \sum_{j=1}^N a_j \frac{\partial u}{\partial x_j} + a_0 u$$

we denote a strongly uniformly elliptic linear differential expression of second order with real-valued coefficient functions $a_{jk} = a_{kj}$, a_j , $a_0 \geq 0$ belonging to $C^\theta(\bar{\Omega})$ ($0 < \theta \leq 1$); m and h are real-valued functions in $C(\bar{\Omega})$, and $\lambda \in \mathbb{R}$ a parameter. In particular, m may change sign in Ω . Let L_0 be the differential operator induced by \mathcal{L} and the Dirichlet boundary conditions, with domain $D(L_0) = \{v \in C^{2+\theta}(\bar{\Omega}) : v = 0 \text{ on } \partial\Omega\}$. Note that L_0 is closable in $C(\bar{\Omega})$ (it admits a closed extension in $L^p(\Omega)$, $1 < p < \infty$, having domain $W_0^{1,p}(\Omega) \cap W^{2,p}(\Omega)$). We set $L := \text{closure of } L_0 \text{ in } C(\bar{\Omega})$. Then L is invertible. For the further study of L we introduce the real Banach spaces $E := C_0(\bar{\Omega}) := \{v \in C(\bar{\Omega}) : v = 0 \text{ on } \partial\Omega\}$ and $X := C_0^1(\bar{\Omega}) := \{v \in C^1(\bar{\Omega}) : v = 0 \text{ on } \partial\Omega\}$, with C -norm $\|\cdot\|_E$ and C^1 -norm $\|\cdot\|_X$, respectively. By the L^p -theory for linear elliptic boundary value problems, L^{-1} maps $C(\bar{\Omega})$ compactly into $X \subset E$. Let finally $M: E \rightarrow E \subset C(\bar{\Omega})$ denote the multiplication operator by the function m .

We define u to be a solution of (1) provided

$$Lu - \lambda Mu = h.\tag{2}$$

Suppose first that $m > 0$ on $\bar{\Omega}$, and let the spaces E and X be provided with the natural ordering given by the cones P_E and P_X of (pointwise) nonnegative functions. The standard notations of ordered Banach spaces are used: $w \geq 0$ if $w \in P$, $w > 0$ if $w \in P \setminus \{0\}$ (P = positive cone). By the strong maximum principle, the compact operator $L^{-1}M$ maps positive functions in E into the interior of P_X (and hence into the quasi-interior of P_E [4, p. 241]). Thus $L^{-1}M: E \rightarrow E$ has positive spectral radius $\mu_1 := \text{spr}(L^{-1}M)$ [4, Th. 3.2, p. 270]. The Krein–Rutman theorem [4, Th. 3.2] guarantees that μ_1 is an eigenvalue of $L^{-1}M$; it is the only eigenvalue of $L^{-1}M$ whose associated eigenspace contains a positive function. Moreover the algebraic multiplicity of μ_1 equals 1. One calls $\lambda_1 := 1/\mu_1 > 0$ the principal eigenvalue of the equation

$$(3) \quad Lu = \lambda Mu;$$

by $u_1 > 0$ we denote the principal eigenfunction. It is proved in [3] that if $\hat{\lambda} \in \mathbb{C}$ is an eigenvalue of the problem obtained from (3) by complexification, then $\text{Re } \hat{\lambda} \geq \lambda_1$.

If $\lambda < \lambda_1$, it is an immediate consequence of the positivity of the operator $L^{-1}M: E \rightarrow E$ that if u is a solution of Eq. (2) with $h \geq 0$, then $u \geq 0$ (for $\lambda \leq 0$ this holds by the maximum principle; for $0 < \lambda < \lambda_1$ we note that (2) is equivalent to

$$u = (I - \lambda L^{-1}M)^{-1}(L^{-1}h),$$

and since $L^{-1}h \geq 0$ and $\text{spr}(\lambda L^{-1}M) < 1$, the assertion follows by representing $(I - \lambda L^{-1}M)^{-1}$ as a Neumann series). For $m \equiv 1$ and $\lambda > \lambda_1$, Clément and Peletier [1] prove an interesting anti-maximum principle.

PROPOSITION 1 [1]. *Let $m \equiv 1$ and $h > 0$. Then there exists a number $\delta > 0$ (depending on h) such that $u < 0$ for the solution u of (2) with $\lambda \in (\lambda_1, \lambda_1 + \delta)$. More precisely, $u \leq c_\lambda u_1$, where $c_\lambda \rightarrow -\infty$ like $(\lambda_1 - \lambda)^{-1}$ as $\lambda \searrow \lambda_1$.*

We now admit functions $m \in C(\bar{\Omega})$ which may change sign. In [2] the hypotheses guaranteeing the existence of a positive principal eigenvalue λ_1 for (3) are drastically weakened. We state the main results obtained there.

PROPOSITION 2 [2]. *Suppose $m \in C(\bar{\Omega})$ is positive at some point in Ω . Then (3) admits a positive principal eigenvalue λ_1 characterized by being the unique positive eigenvalue having a positive eigenfunction u_1 . Moreover $u_1 \in \text{Int}(P_X)$, and λ_1 has the following properties:*

(i) if $\hat{\lambda} \in \mathbb{C}$ is an eigenvalue of the problem obtained from (3) by complexification, with $\operatorname{Re} \hat{\lambda} > 0$, then $\operatorname{Re} \hat{\lambda} \geq \lambda_1$;

(ii) $\mu_1 = 1/\lambda_1$ is an eigenvalue of $L^{-1}M: E \rightarrow E$ with algebraic multiplicity 1.

It is further shown that there is no eigenvalue $\hat{\lambda} \in \mathbb{C}$ with $\operatorname{Re} \hat{\lambda} = 0$.

Regarding Eq. (2), we have

PROPOSITION 3 [2]. Suppose $m \in C(\bar{\Omega})$ is positive somewhere in Ω , and (2) holds with $h \geq 0$. Then

(a) if $0 \leq \lambda < \lambda_1$, it follows that $u \geq 0$;

(b) if $\lambda \geq \lambda_1$, $u > 0$ is possible only in case $\lambda = \lambda_1$ and $h = 0$.

The purpose of this note is to prove that Clément–Peletier's anti-maximum principle carries over to this much more general setting.

THEOREM. Suppose $m \in C(\bar{\Omega})$ admits a positive value at some point in Ω , and let $h > 0$ be given in $C(\bar{\Omega})$. Then there exists a constant $\delta = \delta(h) > 0$ such that if u is the solution of (2) with $\lambda \in (\lambda_1, \lambda_1 + \delta)$, then $u < 0$. In fact, $u \leq c_\lambda u_1$, where $c_\lambda \rightarrow -\infty$ like $(\lambda_1 - \lambda)^{-1}$ as $\lambda \searrow \lambda_1$.

Remark. We consider here only the Dirichlet problem since attention is restricted to Dirichlet boundary conditions in [2]. It is, however, possible to prove the results of [2] also for Neumann or regular oblique derivative boundary conditions.

2. PROOF OF THE THEOREM

We first note that Eq. (2) is equivalent to the equation

$$u - \lambda L^{-1}Mu = L^{-1}h \quad (4)$$

in E . The theorem is proved in a sequence of lemmata.

LEMMA 1. The space E admits the topological direct decomposition

$$E = \operatorname{span}[u_1] \oplus R(I - \lambda_1 L^{-1}M) \quad (5)$$

($u_1 > 0$ is the principal eigenfunction associated with λ_1).

Proof. Since $L^{-1}M: E \rightarrow E$ is compact, $I - \lambda_1 L^{-1}M$ is a Fredholm operator in E with index 0. Hence its range $R(I - \lambda_1 L^{-1}M)$ is closed, and $\operatorname{codim} R(I - \lambda_1 L^{-1}M) = \dim N(I - \lambda_1 L^{-1}M) = 1$. It thus suffices to show that $u_1 \notin R(I - \lambda_1 L^{-1}M)$.

Suppose, to the contrary, that $u_1 = (I - \lambda_1 L^{-1}M)w$ for some $w \in E$. Then $0 = (I - \lambda_1 L^{-1}M)^2 w$. Since $\mu_1 = 1/\lambda_1$ is eigenvalue of $L^{-1}M$ with algebraic multiplicity 1, it follows that $0 = (I - \lambda_1 L^{-1}M)w = u_1$, a contradiction. ■

Let P denote the projection operator in E onto $\text{span}[u_1]$ parallel to $R(I - \lambda_1 L^{-1}M)$, and set $Q := I - P$. Writing

$$L^{-1}h = au_1 + g, \quad u = \beta_\lambda u_1 + v_\lambda$$

($\alpha, \beta_\lambda \in \mathbb{R}$; $g, v_\lambda \in R(I - \lambda_1 L^{-1}M)$), Eq. (4):

$$(I - \lambda_1 L^{-1}M)u + (\lambda_1 - \lambda)L^{-1}Mu = L^{-1}h$$

is equivalent to the system

$$(\lambda_1 - \lambda)\lambda_1^{-1}\beta_\lambda u_1 + (\lambda_1 - \lambda)PL^{-1}Mv_\lambda = au_1 \quad (4a)$$

$$(I - \lambda_1 L^{-1}M)v_\lambda + (\lambda_1 - \lambda)QL^{-1}Mv_\lambda = g \quad (4b)$$

obtained according to the decomposition (5).

LEMMA 2. *There exist constants $\varepsilon > 0$ and $c > 0$ such that $|\lambda_1 - \lambda| < \varepsilon$ implies $\|v_\lambda\|_E \leq c$ for the solution $v_\lambda \in R(I - \lambda_1 L^{-1}M)$ of Eq. (4b) to the parameter value λ .*

Proof. Note that $I - \lambda_1 L^{-1}M$ is an isomorphism in the subspace $R := R(I - \lambda_1 L^{-1}M)$ of E . By the stability of bounded invertibility there exists $\varepsilon > 0$ such that the operator $(I - \lambda_1 L^{-1}M) + (\lambda_1 - \lambda)QL^{-1}M$ remains invertible in R provided $|\lambda_1 - \lambda| < \varepsilon$. The assertion thus follows from Eq. (4b). ■

We need an estimate slightly stronger than that given by Lemma 2. Observe that if $v_\lambda \in R(I - \lambda_1 L^{-1}M)$ is solution of Eq. (4b), then $v_\lambda \in X$. In fact, $g = L^{-1}h - au_1 \in X$; further $L^{-1}Mv_\lambda \in X$ and $QL^{-1}Mv_\lambda = (I - P)L^{-1}Mv_\lambda \in X$ since $PL^{-1}Mv_\lambda = d_\lambda u_1 \in X$ ($d_\lambda \in \mathbb{R}$).

LEMMA 3. *The assertion of Lemma 2 holds with the E -norm replaced by the X -norm.*

Proof. By Lemma 2 we know that $\|v_\lambda\|_E \leq c$ for the solution v_λ of (4b) provided $|\lambda_1 - \lambda| < \varepsilon$. Hence $\|L^{-1}Mv_\lambda\|_X \leq c_1$ and $\|QL^{-1}Mv_\lambda\|_X \leq c_1 + |d_\lambda|\|u_1\|_X \leq c_2$, since $|d_\lambda| \leq d$ by the boundedness of $\{PL^{-1}Mv_\lambda\}$ in E . The result follows again from Eq. (4b). ■

Since $u_1 \in \text{Int}(P_X)$ we infer the existence of a constant $\gamma \in \mathbb{R}$ such that $v_\lambda \leq \gamma u_1$ for the solution v_λ of (4b) to values λ with $|\lambda_1 - \lambda| < \varepsilon$. Equation (4a) implies

$$\beta_\lambda = \alpha\lambda_1(\lambda_1 - \lambda)^{-1} - \lambda_1 d_\lambda \quad (\lambda \neq \lambda_1);$$

thus

$$u \leq c_\lambda u_1 \quad (0 < |\lambda_1 - \lambda| < \varepsilon), \quad (6)$$

where $c_\lambda = \beta_\lambda + \gamma \leq \alpha \lambda_1 (\lambda_1 - \lambda)^{-1} + \lambda_1 d + \gamma$.

LEMMA 4. In the decomposition $L^{-1}h = \alpha u_1 + g$ ($g \in R(I - \lambda_1 L^{-1}M)$), $h > 0$ implies $\alpha > 0$.

Proof. In the present generality we are not able to prove Lemma 4 directly. We show that the cases $\alpha < 0$ and $\alpha = 0$ are impossible.

(i) Assume $\alpha < 0$. Then (6) implies that $u \leq c_\lambda u_1$ with $c_\lambda \rightarrow -\infty$ as $\lambda \nearrow \lambda_1$, contradicting Proposition 3(a).

(ii) Suppose $\alpha = 0$, i.e., $L^{-1}h = w - \lambda_1 L^{-1}Mw$ for some $w \in E$. Without loss of generality we may assume $|m| < 1$ on $\bar{\Omega}$. Then

$$h = Lw - \lambda_1 Mw = (L + \lambda_1)w - \lambda_1(M + 1)w,$$

which gives

$$(L + \lambda_1)^{-1}h = w - \lambda_1 K_{\lambda_1} w, \quad (7)$$

where $K_{\lambda_1} := (L + \lambda_1)^{-1}(M + 1): E \rightarrow E$ is now a compact positive operator. Since $\mu_1 u_1 = K_{\lambda_1} u_1$ ($\mu_1 = 1/\lambda_1, u_1 > 0$), we conclude by the Krein–Rutman theorem that μ_1 is also an eigenvalue of $K_{\lambda_1}^*: E^* \rightarrow E^*$ (the Banach space adjoint operator), with positive eigenfunction u_1^* . From (7) we infer that

$$(L + \lambda_1)^{-1}h \in R(I - \lambda_1 K_{\lambda_1}) = N(I - \lambda_1 K_{\lambda_1}^*)^\perp = (\text{span}[u_1^*])^\perp;$$

consequently

$$\langle u_1^*, (L + \lambda_1)^{-1}h \rangle = 0.$$

On the other hand, $(L + \lambda_1)^{-1}h$ is a quasi-interior point of P_E and hence (since $u_1^* > 0$)

$$\langle u_1^*, (L + \lambda_1)^{-1}h \rangle > 0.$$

This contradiction shows that $\alpha \neq 0$. ■

The assertion of the Theorem now follows immediately from (6) with $\alpha > 0$.

REFERENCES

1. PH. CLÉMENT AND L. A. PELETIER, An anti-maximum principle for second-order elliptic operators, *J. Differential Equations* **34** (1979), 218–229.
2. P. HESS AND T. KATO, On some linear and nonlinear eigenvalue problems with an indefinite weight function, *Comm. Partial Differential Equations* **5** (10) (1980), 999–1030.
3. M. H. PROTTER AND H. F. WEINBERGER, On the spectrum of general second order operators, *Bull. Amer. Math. Soc.* **72** (1966), 251–255.
4. H. H. SCHAEFER, “Topological Vector Spaces,” Macmillan Co., New York, 1966.